

Massless scalar fields and infrared divergences in the inflationary brane world

Oriol Pujolàs^{1*} and Takahiro Tanaka^{2†}

¹ *Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan and*

² *Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

We study the quantum effects induced by bulk scalar fields in a model with a de Sitter (dS) brane in a flat bulk (the Vilenkin-IPser-Sikivie model) in more than four dimensions. In ordinary dS space, it is well known that the stress tensor in the dS invariant vacuum for an effectively massless scalar ($m_{\text{eff}}^2 = m^2 + \xi \mathcal{R} = 0$ with $\mathcal{R}$ the Ricci scalar) is infrared divergent except for the minimally coupled case. The usual procedure to tame this divergence is to replace the dS invariant vacuum by the Allen Follaci (AF) vacuum. The resulting stress tensor breaks dS symmetry but is regular. Similarly, in the brane world context, we find that the dS invariant vacuum generates $\langle T_{\mu\nu} \rangle$ divergent everywhere when the lowest lying mode becomes massless except for massless minimal coupling case. A simple extension of the AF vacuum to the present case avoids this global divergence, but $\langle T_{\mu\nu} \rangle$ remains to be divergent along a timelike axis in the bulk. In this case, singularities also appear along the light cone emanating from the origin in the bulk, although they are so mild that $\langle T_{\mu\nu} \rangle$ stays finite except for non-minimal coupling cases in four or six dimensions. We discuss implications of these results for bulk inflaton models. We also study the evolution of the field perturbations in dS brane world. We find that perturbations grow linearly with time on the brane, as in the case of ordinary dS space. In the bulk, they are asymptotically bounded.

YITP-04-38, KUNS-1927

1. INTRODUCTION

The brane world (BW) scenario [1, 2] has been intensively studied in the recent years. Little is known yet concerning the quantum effects from bulk fields in cosmological models [3, 4, 5, 6]. Quite generically, one expects that local quantities like $\langle T_{\mu\nu} \rangle$ or $\langle \phi^2 \rangle$ can be large close to the branes, due to the well known divergences appearing in Casimir energy density computations. This has been confirmed for example in [7, 8] for flat branes. These divergences are of ultraviolet (UV) nature and do not contribute to the force. Hence, they are ignored in Casimir force computations. However, they are relevant to the BW scenario since they may induce large backreaction, and are worth of investigation.

In this article, we shall shed light on another aspect of objects like $\langle T_{\mu\nu} \rangle$ in BW. We shall point out that they can suffer from infrared (IR) divergences as well. These divergences arise when there is a zero mode in the spectrum of bulk fields in brane models of RSII type with dS brane [1, 9]. The situation is analogous to the case in dS space without brane. It is well known that light scalars in dS develop an IR divergence in the dS invariant vacuum. The main purpose of this article is to explore the effects of scalar fields with light modes in a BW cosmological setup of the RSII type [1]. To consider massless limit of scalar field in inflating BW is especially well motivated in the context of ‘bulk inflaton’ models [5, 6, 10, 11, 12], in which the dynamics of a bulk scalar drives inflation on the brane. In the simplest realizations, the brane geometry is close to dS and the bulk scalar is nearly massless.

Let us recall what happens in the usual dS case [13]. For light scalars $m_{\text{eff}} \ll H$ (with H the Hubble constant) in dS, $\langle \phi^2 \rangle$ and $\langle T_{\mu\nu} \rangle$ in the dS invariant vacuum develop a global IR divergence $\sim 1/m_{\text{eff}}^2$. To be precise, this depends on whether the field is minimally coupled or not. What we have in mind is a generic situation in which the effective mass $m_{\text{eff}}^2 = m^2 + \xi \mathcal{R}$ is small, and $\xi \neq 0$. In these cases $\langle T_{\mu\nu} \rangle$ diverges as mentioned. The point is that in the generic massless limit, another vacuum must be chosen to avoid the global IR divergence. This process breaks dS invariance [14], but this shall not really bother us. The simplest choice is the Allen Follaci (AF) vacuum, in which the stress tensor is globally finite and everywhere regular. The massless minimally coupled case is special [15], and it accepts a different treatment which gives finite $\langle T_{\mu\nu} \rangle$ without violating dS invariance.

In the BW scenario [1], the bulk scalar is decomposed into a continuum of KK modes and bound states. Here we consider the case that there is a unique bound state with mass m_d . If m_d is light, $\langle \phi^2 \rangle$ and $\langle T_{\mu\nu} \rangle$ for the dS invariant vacuum will also diverge like $1/m_d^2$. In this case, again, one will be forced to take another vacuum state like the AF vacuum. Then one naive question is what is the behavior of the stress tensor in such a vacuum in the BW. Also, one might expect singularities on the light cone emanating from the center (the fixed point under the action of dS group)

* pujolas@yukawa.kyoto-u.ac.jp

† tama@scphys.kyoto-u.ac.jp

if we recall that the field perturbations for a massless scalar in dS grow like $\langle\phi^2\rangle \sim \chi$, where χ is the proper time in dS [16, 17, 18, 19] (see also [20]). The light cone in the RSII model corresponds to $\chi \rightarrow \infty$.

Before we start our discussion, we should mention previous calculation given in Ref. [21]. In that paper the stress tensor for a massless minimally coupled scalar was obtained in four dimensions, in the context of open inflation. Montes showed that $\langle T_{\mu\nu} \rangle$ can be regular everywhere except on the bubble. As we will see, these properties hold as well in other dimensions, but only for massless minimal coupling fields.

For simplicity, we consider one extremal case of the RSII model [1] in which the bulk curvature and hence the bulk cosmological constant is negligible. We take into account the gravitational field of the brane by imposing Israel's matching conditions. The resulting spacetime can be constructed by the 'cut-and-paste' technique. Imposing mirror symmetry, one cuts the interior of a dS brane in Minkowski and pastes it to a copy of itself (see Fig. 1). Such a model was introduced in the context of bubble nucleation by Vilenkin [22] and by Ipser and Sikivie [23], and we shall refer to it as 'the VIS model'.

This article is organized as follows. In Section 2, we describe the VIS model and introduce a bulk scalar field with generic bulk and brane couplings. The Green's function is obtained first for the case when the bound state is massive, $m_d > 0$. The form of $\langle T_{\mu\nu} \rangle$ in the limit $m_d \rightarrow 0$ is also obtained. In Section 3, we consider an exactly massless bound state $m_d = 0$, and we present the divergences of the AF vacuum. The case when the bulk mass vanishes is technically simpler and explicit expressions for $\langle T_{\mu\nu} \rangle$ can be obtained. This is done in Section 4. With this, we describe the evolution of the field perturbations in Section 5, and conclude in Section 6.




FIG. 1: (a) Conformal diagram of the Minkowski space. The surfaces corresponding to constant r, χ, τ and ψ coordinates are indicated. The light cone $r = 0$ corresponds to the dashed lines. (b) Conformal diagram of the VIS model.

2. SCALAR FIELDS IN THE VIS MODEL

In this Section we consider a generic scalar field propagating in the VIS model, describing a gravitating brane in an otherwise flat space [22, 23]. Specifically, the space time consists of two copies of the interior of the brane glued at the brane location, as illustrated in Fig. 1.

In the usual Minkowski spherical coordinates the metric is $ds^2 = -dT^2 + dR^2 + R^2 d\Omega_{(n)}^2$, where $d\Omega_{(n)}^2$ stands for the line element on a unit n sphere. From the symmetry, it is convenient to introduce another set of coordinates. The Rindler coordinates, defined by

$$\begin{aligned} R &= r \cosh \chi, \\ T &= r \sinh \chi, \end{aligned}$$

cover the exterior of the light cone emanating from $R = T = 0$. In terms of them the brane location is simply $r = r_0$, and the metric looks like

$$ds^2 = dr^2 + r^2 dS_{(n+1)}^2, \quad (1)$$

where $dS_{(n+1)}^2$ is the line element of de Sitter (dS) space of unit curvature radius. Thus, the Hubble constant on the brane is $H = 1/r_0$. In order to cover the interior of the light cone, we introduce the ‘Milne’ coordinates according to

$$\begin{aligned} R &= \tau \sinh \psi, \\ T &= \tau \cosh \psi. \end{aligned}$$

In these coordinates, the metric is $ds^2 = -d\tau^2 + \tau^2[d\psi^2 + \sinh^2 \psi d\Omega_{(n)}^2]$. Note that we can go from the Rindler to the expanding (contracting) Milne regions making the continuation $r = \pm i\tau$ and $\chi = \psi \mp (\pm)i\pi/2$. Here upper and lower signatures correspond to $+i\epsilon$ and $-i\epsilon$ prescriptions, respectively.

We consider a scalar field even under Z_2 symmetry, with generic couplings described by the action

$$S = -\frac{1}{2} \int_{\text{Bulk}} [(\partial\phi)^2 + (M^2 + \xi R) \phi^2] - \int_{\text{Brane}} [\mu + 2\xi \text{tr} \mathcal{K}] \phi^2, \quad (2)$$

where M and μ are the bulk and brane masses, $\mathcal{R}$ is the Ricci scalar and $\text{tr} \mathcal{K}$ is the trace of the extrinsic curvature. The latter arises because the curvature scalar contains δ function contributions on the brane, and the factor 2 in front of it is due to the Z_2 symmetry. The stress tensor for a classical field configuration can also be split into bulk and a surface parts as [13, 24]

$$\begin{aligned} T_{\mu\nu}^{\text{bulk}} &= (1 - 2\xi) \partial_\mu \phi \partial_\nu \phi - \frac{1 - 4\xi}{2} [(\partial\phi)^2 + (M^2 + \xi \mathcal{R}) \phi^2] g_{\mu\nu} - 2\xi \mathcal{R}_{\mu\nu} \phi^2 - 2\xi \phi \nabla_\mu \nabla_\nu \phi \\ T_{ij}^{\text{brane}} &= [(4\xi - 1) \mu_{\text{eff}} h_{ij} - 2\xi \mathcal{K}_{ij}] \phi^2 \delta(r - r_0), \end{aligned} \quad (3)$$

where h_{ij} is the induced metric on the brane, $\mathcal{R}_{\mu\nu}$ the Ricci tensor, and the equation of motion has been used.¹ Here we have introduced an *effective brane mass* as

$$\mu_{\text{eff}} \equiv \mu + 2(n+1)\xi H, \quad (4)$$

where $H = 1/r_0$ is the Hubble constant on the brane. Then, the v.e.v. $\langle T_{\mu\nu} \rangle$ in point splitting regularization is computed as²

$$\langle T_{\mu\nu} \rangle^{\text{bulk}} = \frac{1}{2} \mathcal{T}_{\mu\nu} [G^{(1)}(x, x')], \quad (5)$$

with

$$\mathcal{T}_{\mu\nu} \equiv \lim_{x' \rightarrow x} \left\{ (1 - 2\xi) \partial_\mu \partial'_\nu - \frac{1 - 4\xi}{2} g_{\mu\nu} (g^{\lambda\sigma} \partial_\lambda \partial'_\sigma + M^2) - 2\xi \nabla_\mu \nabla_\nu \right\}, \quad (6)$$

and

$$\langle T_{ij} \rangle^{\text{brane}} = \frac{1}{2} [(4\xi - 1) \mu_{\text{eff}} h_{ij} - 2\xi \mathcal{K}_{ij}] \delta(r - r_0) G^{(1)}(x, x')|_{x'=x}, \quad (7)$$

where $\partial'_\mu = \partial/\partial x'^\mu$. This expression is extended to the case with a nonzero bulk cosmological constant by replacing M^2 with $M^2 + \xi \mathcal{R}$ and recovering the Ricci tensor term in Eq. (3).

¹ Generically, surface terms are irrelevant for the Casimir force, but are essential to relate the vacuum energy density and the Casimir energy, see [7, 24, 25, 26, 27].

² We omit the anomaly term since it vanishes in the bulk for odd dimension, and on the brane it can be absorbed in a renormalization of the brane tension.

2.1. Spectrum

The Klein-Gordon equation following from the action (2) is separable into radial and dS parts so we introduce the mode decomposition $\phi = \sum \int \mathcal{U}_p(r) \mathcal{Y}_{p\ell m}(\chi, \Omega)$, where m is a multiple index. The radial equation is

$$\left[\partial_r^2 + \frac{n+1}{r} \partial_r + \frac{1}{r^2} \left(p^2 + \frac{n^2}{4} \right) - M^2 \right] \mathcal{U}_p(r) = 0 , \quad (8)$$

while the brane terms can be encoded in the boundary condition

$$(\partial_r + \mu_{\text{eff}}) \mathcal{U}_p|_{r=r_0} = 0, \quad (9)$$

where Z_2 symmetry has been imposed and the effective brane mass μ_{eff} is given in Eq. (4).

The de Sitter part satisfies

$$\left[\square_{n+1} - (n/2)^2 - p^2 \right] \mathcal{Y}_{p\ell m} = 0 .$$

Thus one obtains a tower of modes $\mathcal{Y}_{p\ell m}$ in dS with masses $m_{\text{KK}}^2 = (n/2)^2 + p^2$ in units of H . The mass spectrum determined by the Schrödinger problem defined by Eqs. (8) and (9). It consists of a bound state plus a continuum of KK states with $p \geq 0$ ($m_{\text{KK}} \geq n/2$). The radial part for the KK modes is of the form

$$\mathcal{U}_p^{\text{KK}}(r) = r^{-n/2} [A_p I_{ip}(Mr) + B_p I_{-ip}(Mr)] , \quad (10)$$

with A_p and B_p determined by the boundary condition (9) and continuum normalization, $2 \int dr r^{n-1} \mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_{p'}^{\text{KK}}(r) = \delta(p - p')$.

The mass of the discrete spectrum is

$$m_d^2 = (n/2)^2 + p_d^2 < (n/2)^2 ,$$

and hence p_d is pure imaginary. The normalizability implies that its wave function is

$$\mathcal{U}^{bs}(r) = N_d r^{-n/2} I_{-ip_d}(Mr) , \quad (11)$$

with $-ip_d > 0$. The boundary condition (9) determines p_d in terms of M and μ_{eff} according to

$$\nu I_{-ip_d}(Mr_0) - Mr_0 I'_{-ip_d}(Mr_0) = 0 , \quad (12)$$

where we introduced the combination

$$\nu \equiv \frac{n}{2} - \frac{\mu_{\text{eff}}}{2H} . \quad (13)$$

In the limit $Mr_0 \ll 1$ and $\mu_{\text{eff}} r_0 \ll 1$, Eq. (12) implies that the mass of the bound state is

$$(Hm_d)^2 = nH \mu_{\text{eff}} + \frac{n}{n+2} M^2 + \mathcal{O}(\mu_{\text{eff}}^2, \mu_{\text{eff}} M^2, M^4) , \quad (14)$$




FIG. 2: The shaded area corresponds to the values of μ_{eff} and M^2 for which the bound state is normalizable ($m_d < n/2$) and non tachyonic, $m_d \geq 0$. The thick (red) line corresponds to the massless case. The plot is for $n = 3$.

which agrees with the results of Ref. [10, 28]. Figure 2 shows the values of M^2 and μ_{eff} for which there exists a non-tachyonic ($-ip_d \leq n/2$) bound state. In this paper, we are mostly interested in the situation when the bound state is massless. This happens whenever Eq. (12) with $-ip_d$ replaced by $n/2$ holds, that is when ν reaches the ‘critical’ value

$$\nu_c = \frac{Mr_0 I'_{n/2}(Mr_0)}{I_{n/2}(Mr_0)} . \quad (15)$$

2.2. Green’s function

The renormalized D -dimensional Green’s function can be split into the bound state and KK contributions,

$$G_{(\text{ren})}^{(1)} = G^{\text{KK}} + G^{bs} , \quad (16)$$

with

$$G^{bs} = \mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r') G_{p_d(dS)}^{(1)} , \quad (17)$$

$$G^{\text{KK}} = \int_0^\infty dp [\mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_p^{\text{KK}}(r')]^{ren} G_{p(dS)}^{(1)} , \quad (18)$$

where $G_{p(dS)}^{(1)}$ denotes the Green’s function of a field with mass $(n/2)^2 + p^2$ in $n+1$ dimensional dS space with $H=1$. It depends on x and x' through the invariant distance in dS, which we call ζ . Its precise form is given in Appendix A. The ‘renormalized’ product of the KK mode functions in Eq. (18) is

$$\begin{aligned} [\mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_p^{\text{KK}}(r')]^{ren} &\equiv \mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_p^{\text{KK}}(r') - \mathcal{U}_p^{Mink}(r) \mathcal{U}_p^{Mink}(r') \\ &= \frac{ip}{2\pi(r r')^{n/2}} \frac{\nu K_{-ip}(Mr_0) - Mr_0 K'_{-ip}(Mr_0)}{\nu I_{-ip}(Mr_0) - Mr_0 I'_{-ip}(Mr_0)} I_{-ip}(Mr) I_{-ip}(Mr') + (p \rightarrow -p) . \end{aligned} \quad (19)$$

Here, $\mathcal{U}_p^{Mink}(r) \propto K_{ip}(Mr)$ is the Minkowski counter part of (10). This effectively removes the UV divergent contribution to the Green’s function and guarantees the renormalized Green’s function (16) to be finite in the coincidence limit.

Since Eq. (19) is even in p , Eq. (18) can be cast as $G^{\text{KK}} = \int_{-\infty}^\infty dp [\mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_p^{\text{KK}}(r')]_1^{ren} \big|_{p_d} G_{p(dS)}^{(1)}$, where $[\mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_p^{\text{KK}}(r')]_1^{ren}$ stands for the first term in Eq. (19) only. This can be evaluated summing the residues. From Eq. (A5), the poles in $G_{p(dS)}^{(1)}$ in the upper p plane are at $p = i(q + n/2)$, with $q = 0, 1, 2, \dots$ (see Fig. 3). From Eqs. (19) and (12), we see that the KK radial part has a pole at the value of p corresponding to the bound state, $p = p_d$. We shall now show that the residue is related to the bound state wave function as

$$2\pi i \text{Res} [\mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_p^{\text{KK}}(r')]_1^{ren} \big|_{p_d} = -\mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r') . \quad (20)$$

Using the Wronskian relation $K_\lambda(z) I'_\lambda(z) - K'_\lambda(z) I_\lambda(z) = 1/z$ and Eq. (12), it is straightforward to show that

$$2\pi i \text{Res} [\mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_p^{\text{KK}}(r')]_1^{ren} \big|_{p_d} = -\frac{p_d / I_{-ip_d}(Mr_0)}{\partial_p (\nu I_{-ip}(Mr_0) - Mr_0 I'_{-ip}(Mr_0)) \big|_{p=p_d}} \frac{I_{-ip_d}(Mr) I_{-ip_d}(Mr')}{(r r')^{n/2}} .$$

The overall constant in the r.h.s. is nothing but the normalization constant in the bound state wave function (19), up to the sign. Using the Schrödinger equation (8) and integrating by parts, we have

$$(p^2 - p_d^2) \int_0^{r_0} \frac{dr}{r} I_{-ip_d}(Mr) I_{-ip}(Mr) = I_{-ip}(Mr_0) Mr_0 I'_{-ip_d}(Mr_0) - I_{-ip_d}(Mr_0) Mr_0 I'_{-ip}(Mr_0) .$$

Setting $p = p_d$ after differentiation with respect to p , we find

$$\frac{1}{N_d^2} = 2 \int_0^{r_0} \frac{dr}{r} [I_{-ip_d}(Mr)]^2 = \frac{1}{p_d} I_{-ip_d}(Mr_0) \partial_p (\nu I_{-ip}(Mr_0) - Mr_0 I'_{-ip}(Mr_0)) \big|_{p=p_d} , \quad (21)$$

where we used Eq. (12). From this and the form of the bound state wave function (11), it is clear that Eq. (20) holds. Equation (20) implies that no term of the form $\mathcal{U}_{bs}(r) \mathcal{U}^{bs}(r') G_{p_d}^{(dS)}$ survives in the result. This is ‘fortunate’ because close to $r=0$, $\mathcal{U}^{bs} \sim r^{-n/2-ip_d}$ which is divergent. Thus, Eq. (20) guarantees that $\langle T_{\mu\nu} \rangle$ is regular on the light cone.

Since only the poles from $G_p^{(1)}(dS)$ contribute to $G^{(ren)}$, it can be written as the integral over a contour $\mathcal{C}$ that runs above p_d (see Fig. 3). Equation (A5) leads to the expression appropriate to see the coincidence limit,

$$\begin{aligned} G^{(ren)} &= \int_{\mathcal{C}} dp [\mathcal{U}_p^{\text{KK}}(r) \mathcal{U}_p^{\text{KK}}(r')]_1^{ren} G_p^{(1)}(dS) \\ &= -\frac{S_{(n)}^{-1}}{2^{n-1} \Gamma(\frac{n+1}{2})} \sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \frac{\nu K_{n/2+k+j}(Mr_0) - Mr_0 K'_{n/2+k+j}(Mr_0)}{\nu I_{n/2+k+j}(Mr_0) - Mr_0 I'_{n/2+k+j}(Mr_0)} \\ &\quad \frac{(\frac{n}{2} + k + j) I_{n/2+k+j}(Mr) I_{n/2+k+j}(Mr')}{(rr')^{n/2}} \frac{(-1)^k \Gamma(n + 2k + j)}{j! k! \Gamma(\frac{n+1}{2} + k)} \left(\frac{1 - \cos \zeta}{2} \right)^k, \end{aligned} \quad (22)$$

and we remind that ζ is the invariant distance in dS. Each term comes from the pole at $p = i((k + j) + n/2)$. Setting $k = 0$, we find that the terms with a large j is unsuppressed for $r = r' = r_0$ for $\zeta = 0$. Hence the Green's function in the coincidence limit is divergent on the brane. This is the usual UV 'Casimir' divergence near the boundary. Since we are interested in the IR behavior, we shall not further comment on this UV divergence.

The term with $k = j = 0$ in Eq. (22) renders the Green's function *globally* IR divergent in the limit when the bound state is massless, $\nu \rightarrow \nu_c$ (see Eq. (15)). One can show that this term comes from the homogeneous ($\ell = 0$) mode of the bound state. Using Eqs. (C2) and (C3), the leading behavior of Eq. (22) in the massless limit $m_d \rightarrow 0$ can be written as

$$G^{(ren)(1)} = \frac{2}{S_{(n+1)}} \frac{H^2}{m_d^2} \mathcal{U}_0^{bs}(r) \mathcal{U}_0^{bs}(r') + \mathcal{O}(m_d^0), \quad (23)$$

where $\mathcal{U}_0^{bs}(r) = N_0 I_{n/2}(Mr)/r^{n/2}$ is the wave function of the bound state (11) for the exactly massless case.

The divergence (23) appears because in the massless limit, the wave function of the homogeneous mode of the bound state in the dS invariant vacuum broadens without bound [15]. It can be removed by considering another vacuum with finite width, which implies breaking dS symmetry [14]. Later, we shall take the Allen Follaci vacuum [14]. We will find in Section 3 that in the brane world context, this process removes the global IR divergence, but a localized singularity within the bulk remains.

Let us now examine the behavior of $\langle T_{\mu\nu} \rangle$ in the massless limit, $m_d \rightarrow 0$. In this limit, the stress tensor is given by

$$\langle T_{\mu\nu} \rangle^{bulk} \simeq \frac{H^2}{S_{(n+1)} m_d^2} \mathcal{T}_{\mu\nu} [\mathcal{U}_0^{bs}(r) \mathcal{U}_0^{bs}(r')], \quad (24)$$

$$\langle T_{ij} \rangle^{brane} \simeq \frac{H^2}{S_{(n+1)} m_d^2} [(4\xi - 1)\mu_{\text{eff}} - 2H\xi] (\mathcal{U}_0^{bs}|_{r_0})^2 \delta(r - r_0) h_{ij}, \quad (25)$$

where $\mathcal{T}_{\mu\nu}$ is the differential operator given in Eq. (6) and h_{ij} is the induced metric on the brane. For $M \neq 0$, $\mathcal{U}_0^{bs}$ is not constant. Therefore Eq. (24) explicitly shows the presence of a global IR divergence in $\langle T_{\mu\nu} \rangle$ for the dS invariant vacuum in the $m_d \rightarrow 0$ limit. For $M = 0$, $\mathcal{U}_0^{bs}$ is constant. Hence, the bulk part is finite. However, if $\xi \neq 0$ then the surface term (25) diverges.³ Thus, in the limit $m_d = 0$ we are forced to consider another quantum state. This is the subject of Section 3. We shall mention that there is a possibility to avoid this IR divergence without modifying the choice of dS invariant vacuum state. In the present case the divergence is constant and is localized on the brane. Hence, it can be absorbed by changing the brane tension. We may therefore have a model in which this singular term is appropriately renormalized so as not to diverge in the $m_d \rightarrow 0$ limit. Of course, such a model is a completely different model from the original one without this IR renormalization.

The massless minimally coupled limit, $M = \mu_{\text{eff}} = \xi = 0$, is exceptional. Both bulk and brane parts of stress tensor (24) and (25) are finite in this limit, though here there is a slight subtlety. The limiting values depend on how we fix the ratios among M , μ_{eff} and ξ . For example, using Eq. (14), the surface term is given by

$$\lim_{M, \mu_{\text{eff}}, \xi \rightarrow 0} \langle T_{ij} \rangle^{brane} \simeq -\frac{n+2}{n} \frac{H^2 (\mathcal{U}_0^{bs}|_{r_0})^2}{S_{(n+1)}} \frac{\mu_{\text{eff}} + 2\xi H}{(n+2)H\mu_{\text{eff}} + M^2} \delta(r - r_0), \quad (26)$$

where we used the approximate mass of the bound state,

$$m_d^2 \simeq n\mu H + 2n(n+1)\xi H^2 + \frac{n}{n+2} M^2,$$

³ For $M = 0$, $\mu_{\text{eff}} \langle \phi^2 \rangle$ is finite because $\langle \phi^2 \rangle \sim 1/\mu_{\text{eff}}$, see (Eq. 14).




FIG. 3: Integration contour in Eq. (22).

which is valid in the massless minimal coupling limit (see Eq. (14)). Hence, in the absence of any fine tuning ($m_d^2 \approx \max(M^2, \xi H^2, H\mu)$), it is clear that the contribution (25) is not large, even though the Green's function (23) is. Thus, only in the case when the parameters are 'fine tuned' according to Eq. (15) the stress tensor (25) is large. From Eq. (3), if the Green's function is free from IR divergence, it is clear that the brane stress tensor must be zero in the massless minimally coupled case. The direction that reproduces this result is the one along which $\langle T_{ij} \rangle^{brane}$ already vanishes (in the massive case), that is $\mu_{\text{eff}} = -2\xi H$. Note that this feature is analogous to what happens in dS space [15].

The bulk stress tensor (24) also has a similar but a bit more complicated feature. The operator (6) has terms which do not manifestly involve a small quantity, such as M^2 , ξ or μ . However, these terms are associated with derivative operators. In the limit $M^2 \rightarrow 0$, we can expand the r -dependence of the term with $k = j = 0$ in Eq. (22) as

$$\frac{I_{n/2}(Mr)}{r^{n/2}} \approx \frac{1}{2^{n/2}\Gamma(\frac{n}{2} + 1)} \left(1 + \frac{M^2 r^2}{2(n+2)} + \dots \right). \quad (27)$$

Then, the leading term in the above expansion, which is not suppressed by a factor M^2 , vanishes in (24). The remaining terms are finite unless the parameters are fine tuned, as in the case of the brane stress tensor.

Finally, Eq. (24) also shows that $\langle T_{\mu\nu} \rangle$ is perfectly regular on the light cone for $m_d \neq 0$. As mentioned before, this happens thanks to the KK modes. Note as well that for $M \gtrsim H$, Eq. (24) is exponentially localized on the brane, because the bound state is localized in this case.

3. EXACTLY MASSLESS BOUND STATE

In the preceding section, we have seen that de Sitter (dS) invariant vacuum causes divergence in the limit when the bound state is massless. The divergence is caused by the $\ell = 0$ homogeneous mode in the bound state. In this section we consider a different choice of vacuum state for this mode aiming at resolving the problem of divergence, following the standard methods used in dS space[14, 15]. For simplicity, we concentrate on the case of exactly massless bound state $m_d = 0$ that is $p_d = ni/2$, although more general cases would be treated in a similar way. The case with $m_d = 0$ includes not only a massless minimally coupled scalar, $M = \mu = \xi = 0$, but also other fine tuned cases.

3.1. The Green's function

Here, we should split the Green's function into KK and bound state contributions, $G^{(ren)(1)} = G^{KK(1)} + G^{bs(1)}$. We leave the quantum state for the KK contribution untouched, and change only the contribution from the bound state. In the integral representation for Green's function in (22), we have used the integration contour given in Fig. 3. This choice of contour automatically takes into account the bound state contribution simultaneously. In the present case, we consider the contour that runs below the pole at $p = p_d$ to exclude the bound state contribution. For $m_d = 0$, the integrand has a double pole at $p = ni/2$ because the pole in the radial modes coincides with one of the poles in the dS Green's function. Hence the integral with the contour given in Fig. 3, which runs through these merging poles, is not well defined. But the integral with the new contour, which picks up the contribution from the KK modes only, is well behaved, and it can be cast as

$$G^{KK(1)} = 2\pi i \left\{ \sum_{\text{simple poles}} \text{Res} \left([\mathcal{U}_p^{KK}(r) \mathcal{U}_p^{KK}(r')]_1^{ren} G_p^{(dS)(1)} \right) + \text{Res} \left([\mathcal{U}_p^{KK}(r) \mathcal{U}_p^{KK}(r')]_1^{ren} \right)_{p_d} \partial_p \left[(p - p_d) G_p^{(dS)(1)} \right] \Big|_{p_d} \right. \\ \left. + \text{Res} \left(G_p^{(dS)(1)} \right)_{p_d} \partial_p \left((p - p_d) [\mathcal{U}_p^{KK}(r) \mathcal{U}_p^{KK}(r')]_1^{ren} \right) \Big|_{p_d} \right\}. \quad (28)$$

As before, $[\mathcal{U}_p^{KK}(r) \mathcal{U}_p^{KK}(r')]_1^{ren}$ denotes the first term in Eq. (19). In the first term the ‘simple poles’ mean poles at $p = i(q + n/2)$ with $q = 1, 2, \dots$ (see Fig. 3). Namely, it is obtained by removing the term with $k = j = 0$ from Eq. (22). The last two terms are contributions from the double pole at $p = ni/2$.

Next, we consider the contribution from bound state. A massless bound state behaves as a massless scalar from the viewpoint of $n + 1$ dimensional dS space. In dS space it is well known that the dS invariant Green's function diverges in the massless limit because of the $\ell = 0$ homogeneous mode [14, 15]. The usual procedure is to treat separately the $\ell = 0$ mode from the rest. It is easy to show that (see Appendix B)

$$G^{bs(1)} = \mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r') \left\{ \sum_{\ell > 0} \mathcal{Y}_{\ell m}(\chi) \mathcal{Y}_{\ell m}^*(\chi') + \tilde{\mathcal{Y}}_{AF}(\chi) \tilde{\mathcal{Y}}_{AF}^*(\chi') \right\} + \text{c.c.} \quad (29) \\ = \mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r') \left\{ \partial_p \left[(p - p_d) G_p^{(dS)(+)} \right] \Big|_{p_d} - \partial_p \left[(p - p_d) \mathcal{Y}_{p00}(\chi) \mathcal{Y}_{p00}^*(\chi') \right] \Big|_{p_d} + \tilde{\mathcal{Y}}_{AF}(\chi) \tilde{\mathcal{Y}}_{AF}^*(\chi') \right\} + \text{c.c.},$$

where $\mathcal{Y}_{\ell m}(\chi)$ are the positive frequency dS invariant vacuum modes with mass $p^2 + (n/2)^2$. The last term in Eq. (29) is the contribution from the homogeneous mode in the appropriate state, which we shall take to be the Allen Follaci (AF) vacuum [14] (see Eq. (36)).

At this stage, we note that the second term of Eq. (28) cancels the first term of Eq. (29), due to Eq. (20). This cancelation resembles the one that occurred in the previous case between the KK modes and the bound state. In that case, it guaranteed the absence of light cone divergences. In the present case, the terms that cancel in Eqs. (28) and (29) are already regular. We show below (see discussion after Eq. (32)) that instead the last term in Eq. (28) and the second term in Eq. (29) diverge on the light cone. However, when added up, they render G^{ren} finite on the light cone in odd dimension. (In even dimensions, G^{ren} is finite but its derivatives are not.)

The fact that the dS invariant Green's function diverges because of the homogeneous mode implies that $\text{Res}(G_p^{(dS)(1)})_{p_d} = \lim_{p \rightarrow p_d} (p - p_d) \mathcal{Y}_{p00} \mathcal{Y}_{p00}^* + \text{c.c.}$. Using this and Eq. (20), we can rewrite the total Green's function in the more convenient form

$$G^{(ren)(1)} = G_{simple}^{(1)} + G_{LC} + \tilde{G}_{AF}, \quad (30)$$

with

$$G_{LC} \equiv 2\pi i \partial_p \left((p - p_d) \mathcal{Y}_{p00} \mathcal{Y}_{p00}^* - (p - p_d) [\mathcal{U}_p^{KK}(r) \mathcal{U}_p^{KK}(r')]_1^{ren} \right) \Big|_{p_d} + \text{c.c.}, \\ \tilde{G}_{AF} \equiv \mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r') \tilde{\mathcal{Y}}_{AF} \tilde{\mathcal{Y}}_{AF}^* + \text{c.c.}, \quad (31)$$

and $G_{simple}^{(1)}$ is the contribution from the first term in Eq. (28). As is manifest from the expression (22) with the $k = j = 0$ term removed, $G_{simple}^{(1)}$ is regular in the coincidence limit, everywhere except for the ‘Casimir’ divergences on the brane, and depends on the points x and x' through r , r' and $\zeta(x, x')$ (the invariant distance in dS between the projections of points x and x'), and hence it is dS invariant. The second term, G_{LC} , contains the two contributions in Eqs. (28) and (29) that are separately divergent on the light cone mentioned in the previous paragraph. The last term, $\tilde{G}_{AF}$, encodes the choice of vacuum for the zero mode of the bound state.

Let us examine the contribution potentially divergent on the light cone G_{LC} . The derivative of the χ dependent part is obtained from Eqs. (B4) and (B6). The derivative of the radial part follows from Eq. (19) and $\partial_\lambda I_\lambda(z) = I_\lambda(z) \log z + f(z)$, where $f(z)$ is a regular function. It is easy to see that G_{LC} takes the form

$$G_{LC}(x, x') = \frac{2}{nS_{(n+1)}} \mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r') \operatorname{Re} [F(x) + F(x')] + \text{regular}, \quad (32)$$

with

$$F(x) = \log r + n \int_0^\chi \frac{d\chi_1}{\cosh^n \chi_1} \int_{-\frac{i\pi}{2}}^{\chi_1} d\chi_2 \cosh^n \chi_2, \quad (33)$$

where the indicated regular term depends on r and r' only. The double integral in (33) grows linearly with χ for large χ and eventually blows up on the light cone (The first integral asymptotically grows like $e^{n\chi_1}/n$, and therefore the integrand in the second integral goes to a constant). This is the expected behavior from the massless bound state. On the other hand, the KK modes contribute the $\log r$ term, which cancels the light cone divergence. To see this, we integrate by parts to obtain

$$F(x) = \log r + \log \sinh \chi + \int_0^\chi \frac{d\chi_1}{\cosh^n \chi_1} \int_{-\frac{i\pi}{2}}^{\chi_1} d\chi_2 \frac{\cosh^n \chi_2}{\sinh^2 \chi_2}, \quad (34)$$

and now the double integral is bounded. The first two terms are simply $\log T$. Thus, the leading divergence in G_{LC} on the light cone cancels between contributions from bound state and from the KK modes, although it still diverges logarithmically at infinity.

This statement has to be qualified for even dimension. In this case, the derivatives of G_{LC} diverge on the light cone because of the last term in Eq. (34). To see this, note that the integrand in Eq. (34) can be expanded in exponentials. Then, the integral is a sum of exponential terms except for one, of the form $\chi e^{-n\chi}$, if n is even. In terms of the null coordinates U and V , this is $\sim (V/U)^{n/2} \log(V/U)$ for $\chi \rightarrow +\infty$ (for $\chi \rightarrow -\infty$ replace $U \leftrightarrow V$). Even though the Green's function is regular at $V = 0$, the stress tensor develops a singularity which behaves like $\sim 1/V$ or $1/U$ on the light cone in four dimensions ($n = 2$) and like $\sim \log V$ or $\log U$ in six dimensions ($n = 4$) if $\xi \neq 0$.

For reference, we show the explicit form of $F(x)$ for dimensions 4 and 5,

$$F(x) = \begin{cases} \frac{U \log U + V \log |V|}{R} & \text{for } n = 2, \\ \log(T + \tau) - \frac{(T - \tau)\tau}{R^2} & \text{for } n = 3, \end{cases} \quad (35)$$

where $\tau^2 = T^2 - R^2$. Note that despite appearances, Eq. (35) is regular at $R = 0$. This is guaranteed since the χ dependent part of $F(x)$ is related to the dS invariant vacuum modes (see Eq. (B6)), which are regular at $R = 0$ ($\xi = -\pi i/2$) by construction. The expressions (35) are appropriate in the Milne region, and we have already taken the real part, which is the relevant part for $G^{(1)}$.

The last term $\tilde{G}_{AF}$ in Eq. (30) corresponds to the choice of vacuum for the $\ell = 0$ mode. This mode is peculiar because it behaves like a free particle rather than an oscillator. The eigenstates of the Hamiltonian, and in particular the ground state, are plane waves in field space. However, such states are not normalizable. One can construct well defined states as wave packets. The simplest option is a Gaussian packet. This is the Allen Follaci vacuum [14], which in fact is a two-parameter family of vacua. Its mode function is given by [14]

$$\tilde{\mathcal{Y}}_{AF}(\chi) = \frac{1}{\sqrt{S_{(n)}}} \left[\frac{1}{2\alpha} + i\beta - i\alpha \int_0^\chi \frac{d\chi'}{\cosh^n \chi'} \right], \quad (36)$$

where $\alpha > 0$ and β are the mentioned free (real) parameters. We shall impose that the vacuum is time reversal symmetric. This translates into $\tilde{\mathcal{Y}}_{AF}(-\chi) = \tilde{\mathcal{Y}}_{AF}^*(\chi)$, and implies $\beta = 0$. Because of the time dependence, it breaks dS symmetry. For this vacuum,

$$\tilde{G}_{AF}(x, x') = \frac{\mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r')}{S_{(n)}} \left\{ \frac{1}{2\alpha^2} + 2\alpha^2 \operatorname{Re} [F_{AF}(x) F_{AF}^*(x')] - \operatorname{Im} [F_{AF}(x) + F_{AF}(x')] \right\} \quad (37)$$

with

$$F_{AF} = \int d\chi \cosh^{-n} \chi.$$

In order to obtain the analytic continuation of F_{AF} to the Milne region, it is better to write it as an integral over a contour with constant T . It is straightforward to see that

$$F_{AF}(x) = -T \int^R \frac{dR'}{R'^n} (R'^2 - T^2)^{\frac{n}{2}-1}.$$

It is transparent now that F_{AF} behaves like $1/R^{n-1}$ near $R = 0$. It is also clear that it is regular on the light cone and is bounded at infinity. Moreover, it gets an imaginary part in the Milne region for odd dimensions (in the Rindler region it is always real). More specifically, we have

$$F_{AF}(x) = \begin{cases} \frac{T}{R}, & \text{for } n = 2, \\ \frac{i}{2} \left[\frac{T\tau}{R^2} - \ln \left(i \frac{T+\tau}{R} \right) \right] & \text{for } n = 3. \end{cases} \quad (38)$$

3.2. Divergent stress tensor

Now we discuss the form of the expectation value of the stress tensor for the possible values of M , μ_{eff} and ξ in which the bound state is massless. The bulk part of the stress tensor is most conveniently separated into

$$\langle T_{\mu\nu} \rangle = \langle T_{\mu\nu} \rangle_0 + \langle T_{\mu\nu} \rangle_{\text{simple}} \quad (39)$$

where $\langle T_{\mu\nu} \rangle_0$ contains the contributions from G_{LC} and $\tilde{G}_{AF}$, whereas $\langle T_{\mu\nu} \rangle_{\text{simple}}$ is the contribution from $G_{\text{simple}}^{(1)}$. All the IR irregularities are contained in $\langle T_{\mu\nu} \rangle_0^{\text{bulk}}$. From Eqs. (32) and (37), we obtain the bulk part of $\langle T_{\mu\nu} \rangle_0$ as

$$\begin{aligned} \langle T_{\mu\nu} \rangle_0^{\text{bulk}} &= \frac{1}{2S_{(n)}} \mathcal{T}_{\mu\nu} \left[\mathcal{U}_{bs}(r) \mathcal{U}_{bs}(r') \left\{ \frac{1}{2\alpha^2} + 2\alpha^2 \text{Re}[F_{AF}(x) F_{AF}^*(x')] \right\} \right] \\ &\quad - 2\xi \nabla_\mu \nabla_\nu \mathcal{U}_{bs}(r) \mathcal{U}_{bs}(r') \left[\frac{\text{Re } F(x)}{nS_{(n+1)}} - \frac{\text{Im } F_{AF}(x)}{2S_{(n)}} \right]. \end{aligned} \quad (40)$$

The term proportional to α^2 diverges at $R = 0$ like $1/R^{2n}$ for any value of ξ . $R = 0$ corresponds to a timelike axis in the bulk passing through the center of symmetry (see Fig. 5). The last term also diverges like $1/R^{n+1}$ for odd dimensions, but vanishes for even dimensions. The piece involving $\text{Re } F$ diverges on the light cone for $n = 2$ or 4 .

Let us begin with the most general case with $M \neq 0$. In this case $\mathcal{U}^{bs}(r)$ is not constant, and the contribution from the term inversely proportional to α^2 in Eq. (40) does not vanish. This term is analogous to Eq. (24), and it diverges globally in the $\alpha \rightarrow 0$ limit. Hence, this state cannot be taken, and one has to content with the AF vacuum with some nonzero α . Hence, the term proportional to α^2 is unavoidable. But this is very noticeable since it contains a singularity at $R = 0$ of the form $\sim \alpha^2/R^{2n}$, present even for minimal coupling. The main point is that in the presence of a bulk mass, $\langle T_{\mu\nu} \rangle$ contains a quite severe bulk singularity even after we get rid of the global IR divergence. There is of course the possibility that a different choice of vacua for the KK modes could cancel it out. In this case, it seems that the vacuum choice should not be dS invariant, otherwise the singular zero mode contribution that is not dS invariant could not be compensated. We leave this issue for future research.

Now we turn to the case with $M = 0$. Let us first consider the nonminimal case $\xi \neq 0$. Since $M = 0$, $\mathcal{U}^{bs}(r)$ is constant and the bulk stress tensor does not diverge globally in the $\alpha \rightarrow 0$ limit. Divergences in the other α independent two terms in (40) also vanish in even dimensions with $n \geq 6$. However, we cannot avoid the divergence in the brane stress tensor in the $\alpha \rightarrow 0$ limit. From Eq. (7) and taking into account that $\mu_{\text{eff}} = 0$ (see Eq. (14)), we have

$$\begin{aligned} \langle T_{ij} \rangle_0^{\text{brane}} &= -2\xi \langle \phi^2 \rangle_{AF} \mathcal{K}_{ij} \delta(r - r_0) \\ &= -\frac{\xi n}{r_0^{n+1}} \left(\frac{(2\alpha)^{-2} + \alpha^2 F_{AF}^2(x)}{S_{(n)}} + \frac{2\text{Re } F(x)}{S_{(n+1)}} \right) h_{ij} \delta(r - r_0), \end{aligned} \quad (41)$$

where we used that $\mathcal{U}_{bs}^2 = n/2r_0^n$. Thus, one is forced to take the AF vacuum again. As a result, the bulk stress tensor develops the same singularity at $R = 0$ as before in the $M \neq 0$ case. The globally divergent $(1/\alpha^2)$ term in Eq. (41) is proportional to the induced metric. One might wonder whether this effect is physical or not, since it could be simply absorbed in the brane tension as before. We think that such a procedure is not justified here because α is a state dependent parameter, and renormalization should be done independently of the choice of the quantum state.

Before we examine the stress tensor in the massless minimal coupling case, let us comment on the relation between the AF vacuum and the Garriga Kirsten (GK) vacuum. The latter was introduced in [15] for a massless minimally coupled scalar in dS. It corresponds to the plane wave state with zero momentum in field space. This is intrinsically ill defined, giving a ‘constant infinite’ contribution to the Green’s function. However, the point is that, since it is an eigenstate of the Hamiltonian, it does not depend on χ so it is dS invariant. Divergence in the Green’s function can be accepted in the massless minimally coupled case. Our reasoning is as follows. In this case, the action has the symmetry $\phi \rightarrow \phi + \text{constant}$. If we consider it as a ‘gauge’ symmetry, all the observables are to be constructed from derivatives (or differences) of the field. In fact the stress tensor operator (6) only contains derivatives of the type $\partial_\mu \partial'_\nu G^{(1)}(x, x')$ in this case. Then we will find that the constant contributions in the Green’s function are irrelevant.⁴ We need a practical way to compute quantities in this vacuum. If we follow the argument by Kirsten and Garriga, it will be given by the limit of AF vacuum

$$|0\rangle_{GK} \equiv \lim_{\alpha \rightarrow 0} |0\rangle_{AF} .$$

From Eq. (41), it follows that the brane term $\langle T_{ij} \rangle^{brane}$ vanish in the GK vacuum. The first term in Eq. (40), which is responsible for the ‘infinite constant’, vanishes for $M = 0$. While the second term proportional to α^2 also vanishes in the $\alpha \rightarrow 0$ limit. We shall note that the last two terms in Eq. (40) independent of α are also zero for $\xi = 0$. Hence, we are left with $\langle T_{\mu\nu} \rangle_{simple}$, which is manifestly dS invariant, and is finite (aside from the ‘Casimir’ divergences on the brane). Thus, the total stress tensor in the GK vacuum is given by a simple formula presented below in Eq. (50) with $\xi = 0$. In contrast to the massless minimal coupling limit discussed in the preceding section, here we do not have any ambiguity.

4. ZERO BULK MASS

In the absence of bulk mass M , the Green’s function and the stress tensor can be obtained explicitly. We shall thus discuss now this case.

4.1. Generic mass of the bound state

First we discuss the case when the bound state is not massless. The other case is postponed until Section 4.2.

For the bound state, the boundary condition (9) fixes $p_d = i\nu$ and m_d is given by Eq. (14) with $M = 0$. Its wave function is proportional to $r^{-ip_d - n/2} = r^{-\mu_{eff}/H}$, which is constant if $m_d = 0$ (see Eq. (14)). From the discussion in Section 2.1, the bound state is normalizable and non-tachyonic for $n/2 \geq \nu > 0$ (i.e. $0 \leq \mu_{eff} < (n/2)H$), see Fig. 2.

The normalized radial KK modes are

$$\mathcal{U}_p^{KK}(r) = \sqrt{\frac{2}{\pi r^n (1 + (\nu/p)^2)}} \left(\cos(p \ln r) + \frac{\nu}{p} \sin(p \ln r) \right) , \quad (42)$$

and the renormalized product of mode functions in Eq. (18) is

$$[\mathcal{U}_p^{KK}(r) \mathcal{U}_p^{KK}(r')]^{ren} = \frac{1}{2\pi r^{n/2}} \left[\frac{p + i\nu}{p - i\nu} (rr')^{-ip} + \frac{p - i\nu}{p + i\nu} (rr')^{ip} \right] . \quad (43)$$

Proceeding as in Eq. (22), we can explicitly perform the integration over the KK modes and in this case, we obtain a simpler expression,

$$D_{(ren)}^{(1)}(x, x') = \frac{1}{(2r_0)^n S_{(n)} \Gamma(\frac{n+1}{2})} \sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \frac{\frac{n}{2} + \nu + k + j}{\frac{n}{2} - \nu + k + j} \frac{(-1)^k \Gamma(n + 2k + j)}{j! k! \Gamma(\frac{n+1}{2} + k)} \left(\frac{rr'}{r_0^2} \right)^{k+j} \left(\frac{1 - \cos \zeta}{2} \right)^k , \quad (44)$$

⁴ Taking this vacuum is analogous to performing a Gupta-Bleuler quantization [29].

where as usual, we denote the (bulk-) massless Green's function by $D_{(ren)}^{(1)}(x, x')$. From Eq. (44), it is manifest that the Green's function in the dS invariant vacuum state is regular at $r = 0$ in the coincidence limit. As before, in the massless limit $\nu \rightarrow n/2$, it is singular. However, since the divergent term $k = j = 0$ is constant in the present case, it will not affect the bulk $\langle T_{\mu\nu} \rangle$.

In order to compute the stress tensor, it is convenient to rewrite Eq. (44) in a more compact form. For a conformally coupled field, ν vanishes, and the Green's function actually takes the simple form⁵

$$D_{\nu=0}^{(ren)(1)}(x, x') = \frac{1}{nS_{(n+1)}} \left(\frac{1}{r_0^2 + (rr'/r_0)^2 - 2rr' \cos \zeta} \right)^{n/2}. \quad (45)$$

This expression can be obtained by the method of images. It corresponds to the potential induced by a source of unit charge at x' together with an image source located at $r'_I = r_0^2/r'$ with a charge $q'_I = (r'_I/r_0)^n$. From the form of (44), the Green's function for $\nu \neq 0$ can be obtained from Eq. (45) by applying the integral operator

$$D^{(ren)(1)} = \left[1 + 4\nu r_0^{-2\nu} \int_{r_0}^{\infty} \frac{d\tilde{r}_0}{\tilde{r}_0} \tilde{r}_0^{2\nu} \right] D_{\nu=0}^{(ren)(1)}. \quad (46)$$

Borrowing the intuition from the method of images, Eq. (46) can be interpreted as the potential induced by the image charge mentioned in the previous paragraph together with a string stretching from r_0^2/r' to infinity along the radial direction defined by x' with a charge line density given by $\lambda(r) = 4\nu(r/r_0)^{2\nu-1}/r'$. To obtain the stress tensor, we can first compute the case $\nu = 0$ and then apply the same operator as in Eq. (46). The general result with $M = 0$ and $m_d \neq 0$ is

$$\begin{aligned} {}^{(m_d \neq 0)}\langle T_r^r \rangle &= (\xi - \xi_c) \frac{(-1)^n (n+1)}{S_{(n+1)} r_0^{n+2}} \left\{ \frac{r_0^{2n+2}}{(r_0^2 - r^2)^{n+1}} + \frac{4\nu}{n+2-2\nu} {}_2F_1 \left(n+1, \frac{n}{2} + 1 - \nu; \frac{n}{2} + 2 - \nu; \frac{r^2}{r_0^2} \right) \right\}, \\ {}^{(m_d \neq 0)}\langle T_j^i \rangle &= (\xi - \xi_c) \frac{(-1)^n (n+1)}{S_{(n+1)} r_0^{n+2}} \left\{ \frac{r_0^{2n+2} (r_0^2 + r^2)}{(r_0^2 - r^2)^{n+2}} + \frac{4\nu}{n+2-2\nu} {}_2F_1 \left(n+1, \frac{n}{2} + 1 - \nu; \frac{n}{2} + 2 - \nu; \frac{r^2}{r_0^2} \right) \right. \\ &\quad \left. + \frac{8\nu}{n+4-2\nu} \frac{r^2}{r_0^2} {}_2F_1 \left(n+2, \frac{n}{2} + 2 - \nu; \frac{n}{2} + 3 - \nu; \frac{r^2}{r_0^2} \right) \right\} \delta_j^i, \end{aligned} \quad (47)$$

where $\xi_c = n/4(n+1)$. As an aside, we shall note as well that for conformal coupling, $\langle T_{\mu\nu} \rangle^{bulk} = 0$ even with a nonzero boundary mass μ . This is a consequence of conservation and tracelessness of $T_{\mu\nu}$.

4.2. Massless bound state

We consider now the case $m_d = 0$, that is $\nu = n/2$. We can proceed as in Eq. (30) and decompose $D^{ren} = D_{simple} + G_{LC} + \tilde{G}_{AF}$, where D_{simple} is the contribution from the simple poles $p = i(n/2 + k)$ with $k = 1, 2, \dots$ in Eq. (28). Thus, it is given by the terms in Eq. (44) with non-vanishing k and j . The integral representation analogous to Eq. (46) is now

$$D_{simple}^{(1)} = \left[1 + \frac{2n}{r_0^n} \int_{r_0}^{\infty} \frac{d\tilde{r}_0}{\tilde{r}_0} \tilde{r}_0^n \right] \left(D_{\nu=0}^{(ren)} - \frac{1}{nS_{(n+1)} \tilde{r}_0^n} \right), \quad (48)$$

where we subtract the constant to remove the $j = k = 0$ term. The explicit expression in four dimensions was obtained in [21]⁶, and we shall not reproduce it here. The case of main interest for us is $n = 3$ and we find, up to a finite constant,

$$D_{simple}^{(1)}(x, x') = \text{Re} \left[\frac{1}{8\pi^2} \frac{1}{r_0^3 \Delta^{3/2}} + \frac{3}{8\pi^2 r_0^3} \left\{ \frac{\cos 2\zeta}{\sin^2 \zeta} - \frac{\cos 2\zeta - (rr'/r_0^2) \cos \zeta}{\sin^2 \zeta \Delta^{1/2}} - \log \left[1 - \frac{rr'}{r_0^2} \cos \zeta + \Delta^{1/2} \right] \right\} \right], \quad (49)$$

⁵ From Eq. (44), the expression for a Dirichlet scalar ($\nu \rightarrow \infty$) is the same with opposite sign.

⁶ In Ref. Z_2 reflection symmetry is not assumed for the field.

where $\Delta = 1 + (rr'/r_0^2)^2 - 2(rr'/r_0^2)\cos\zeta$. As mentioned above, in the coincidence limit this contribution is regular except on the brane. We shall note as well that it grows logarithmically at infinity.

The contribution $\langle T_{\mu\nu} \rangle_{simple}^{bulk}$ is easily found exploiting the integral representation (48), as before. The only difference between (48) and (46) in the limit $\nu = n/2$ (*i.e.* $m_d = 0$) is a constant, which does not affect $\langle T_{\mu\nu} \rangle$ in this case. Thus, $\langle T_{\mu\nu} \rangle_{simple}^{bulk}$ reduces to (47) with $\nu = n/2$. This gives a simple expression in terms of elementary functions. In four dimensions it is given in [21] (for $\xi = 0$). In five dimensions, we obtain

$$\begin{aligned} \langle T_r^r \rangle_{simple}^{bulk} &= -\frac{9}{32\pi^2} \frac{\xi - \xi_c}{r_0^3} \left\{ \frac{r_0^6}{(r_0^2 - r^2)^4} + \frac{r^4 - 3r_0^2 r^2 + 3r_0^4}{(r_0^2 - r^2)^3} \right\} , \\ \langle T_j^i \rangle_{simple}^{bulk} &= -\frac{9}{32\pi^2} \frac{\xi - \xi_c}{r_0^3} \left\{ r_0^6 \frac{r^2 + r_0^2}{(r_0^2 - r^2)^5} + \frac{r^2}{2} \frac{r^4 - 4r_0^2 r^2 + 6r_0^4}{(r_0^2 - r^2)^4} + \frac{r^4 - 3r_0^2 r^2 + 3r_0^4}{(r_0^2 - r^2)^3} \right\} \delta^i_j , \end{aligned} \quad (50)$$

which is dS invariant and regular everywhere except on the brane.

5. PERTURBATIONS

We shall now discuss the form of the field perturbations, focusing on the bulk massless case with $n = 3$, since we have obtained a closed form expression for the Green's function and the stress tensor in the preceding section. We begin by the case with generic coupling, and we consider $\langle \phi^2(x) \rangle$ in the Allen Follaci vacuum. The massless minimally coupled case in the de Sitter (dS) invariant Garriga Kirsten vacuum requires a different treatment, which will be discussed in Section 5.2.

5.1. Generic coupling

The renormalized expectation value of $\phi^2(x)$ is given by

$$\langle \phi^2(x) \rangle_{AF}^{ren} = \frac{1}{2} G_{ren}^{(1)}(x, x) ,$$

with $G_{ren}^{(1)}$ given by (30). From Eqs. (37) and (38), it diverges at $R = 0$ because of G_{AF} . It is also clear that this contribution is bounded at (null) infinity. From Eq. (32), the G_{LC} contribution is regular on the light cone. Equation (49) shows that the KK contribution D_{simple} diverges on the brane. In the bulk, both D_{simple} and G_{LC} grow logarithmically at infinity. However, the growing terms cancel out. Indeed, Eq. (49) shows that as long as x is not on the brane, D_{simple} at coincident points is dominated by the logarithmic term. Therefore we have

$$\frac{1}{2} D_{simple}^{(1)}(x, x) \sim -\frac{3}{16\pi^2 r_0^3} \log \left| 1 - \frac{r^2}{r_0^2} \right| , \quad (51)$$

in the limit $x \rightarrow \infty$. As for G_{LC} , since the wave function of the bound state in the $M = 0$ case is $\mathcal{U}_{bs}^2 = n/2r_0^n$, we find

$$\frac{1}{2} G_{LC}(x, x) = \frac{3}{8\pi^2 r_0^3} \text{Re}[\log(T - ir)] + \mathcal{O}(1) , \quad (52)$$

where we used Eq. (35). It is clear that the logarithmic terms cancel and as a result $\langle \phi^2(x) \rangle_{AF}^{ren}$ is bounded at infinity. This is expected, because the bulk is flat. Intuitively, in four dimensional dS space, the perturbations grow because when the modes are stretched to a super-horizon scale, they freeze out. Since modes of ever smaller scales continuously being stretched, they pile up at a constant rate [19]. Since this effect is due to the local curvature of the spacetime, it should not happen in a flat bulk.

Accordingly, on the brane we recover the same behavior as in de Sitter space. Indeed, restricting (52) on the brane, we obtain $G_{LC}^{(1)} \sim \chi$. We have mentioned before that $D_{simple}(x, x)$ is UV divergent on the brane. Since this happens because point splitting regularization used here does not operate on the brane, this object needs UV regularization

and renormalization. This can be done in a variety of ways, *e.g.* with dimensional regularization⁷, introducing a finite brane thickness, smearing the field etc. The point is that the renormalized value must be a constant simply because $D_{simple}(x, x)$ is dS invariant and is a function of x only. Thus, Eqs. (51) and (52) imply that on the brane, $\langle \phi^2 \rangle$ grows in time, as in dS space.

5.2. Massless minimal coupling

Now, we shall see essentially the same features arise for a massless minimally coupled scalar in the GK vacuum, in which case everything can be obtained in a dS invariant way [15]. Because of the shift symmetry $\phi \rightarrow \phi + \text{const.}$, $\langle \phi^2 \rangle$ is not an observable in this case. Still, it is possible to define a ‘shift-invariant’ notion for the field perturbations. Following [15], one introduces the correlator

$$\mathcal{G}(x, y) \equiv \langle [\phi(x) - \phi(y)]^2 \rangle_{GK}, \quad (53)$$

which can be thought of as the combination $[G^{(1)}(x, x) + G^{(1)}(y, y) - 2G^{(1)}(x, y)]/2$. Since the first two terms are UV divergent, we shall rather consider

$$\mathcal{G}^{ren}(x, y) = \frac{1}{2} [G^{ren(1)}(x, x) + G^{ren(1)}(y, y) - 2G^{ren(1)}(x, y)]. \quad (54)$$

The main point is that all the terms of the form $f(x) + f(y)$ in $G^{ren(1)}$ cancel out in the combination $\mathcal{G}^{ren}(x, y)$. All the terms in Eqs. (32) and (37) that do not vanish in the limit $\alpha \rightarrow 0$ are of this form because $\mathcal{U}^{bs}$ is constant in the massless minimally coupled case. Thus, this correlator is well defined for the GK vacuum [15] and we can readily write

$$\mathcal{G}^{ren}(x, y) = \frac{1}{2} [D_{simple}^{(1)}(x, x) + D_{simple}^{(1)}(y, y) - 2D_{simple}^{(1)}(x, y)], \quad (55)$$

with $D_{simple}(x, y)$ given by (49).⁸ Thus, the behavior of the perturbations in the GK vacuum for x and y distant is thus described by the asymptotic behavior of $D_{simple}(x, x)$ and $D_{simple}(x, y)$. The former is summarized in Eq. (51). As mentioned before, for finite x , $D_{simple}(x, x)$ is regular everywhere except on the brane (in which case $y_I = y$).

Before describing the asymptotic form of $D_{simple}(x, y)$, we need to discuss the singularities that it contains. The combination that we called Δ in Eq. (49) is

$$\Delta(x, y) = |x - y_I|^2 / |y_I|^2,$$

where $|x|^2 = \eta_{\mu\nu} x^\mu x^\nu$ and $y_I^\mu = (r_0^2/|y|^2)y^\mu$ is the image of x (see comments around Eq. (46) and Fig. 5). Note that when y is in one of the Milne regions, then y_I is in the other one. Rather than an ‘image charge’, it represents the point where the light cone focuses, (see Fig. 4 (b)). For $x \neq y$ and both finite, $D_{simple}^{(1)}(x, y)$ has singularities in two types of situations. One is when x is on the light cone of y_I (then, $\Delta = 0$). The other case arise when the argument of the logarithmic term in Eq. (49) vanishes. It is convenient to rewrite this term as

$$-\frac{3}{8\pi^2 r_0^3} \log \left| \frac{(|x - y_I| + |y_I|)^2 - |x|^2}{2|y_I|^2} \right|. \quad (56)$$

The argument vanishes when they are aligned with respect to the origin *and* $|x|^2 > |y_I|^2$. This condition defines an ‘image string’ stretching from y_I to infinity. Hence, this singularity occurs only when two points are in different Milne regions. This is consistent with the interpretation of Eq. (46), that the Green’s function can be constructed with a

⁷ It is illustrative to consider the massless conformal case. The Green’s function (45) for x and x' on the brane is simply $D_{\nu=0}^{(ren)}(x, x') \sim 1/|x - x'|^{n/2}$. If n is negative, this is clearly zero in the coincidence limit, and hence the continuation to $n = 3$ is zero as well. In the non-conformal case, we can use the integral form (46) to compute $\langle \phi^2 \rangle \delta(r - r_0)$ in arbitrary dimension. Upon continuation to $n = 3$, one obtains a pole $\sim 1/(n - 3)$ and a nonzero finite part. The pole can be absorbed in the brane tension and $\langle \phi^2(r_0) \rangle^{ren}$ is a finite constant.

⁸ Note that (55) is completely regular on the light cone.




FIG. 4: The singularities in $D_{\text{simple}}(x, y)$ are located on the light cone emanating from y_I , the image point of y . Also when x is on the light wedge emanating from ‘image string’ stretching from y_I to infinity. Here, we plot the image point (smaller filled circle) with the image string attached to it. In (a), y is spacelike, in (b) null, in (c) timelike and in (d) at null infinity.

mirror image and a linear charge distribution over such a string, see Fig. 4.⁹ These two situations correspond to the coincidence of x with the image of y or its ‘light cone’. Therefore all of them are of UV nature.¹⁰

The asymptotic form of $D_{\text{simple}}(x, y)$ for distant points follows from Eq. (56) because then, the inverse powers of Δ in (49) are finite. Taking y fixed and $x \rightarrow \infty$ (with x not on the light cone from y_I nor on the image string nor on its light wedge), one obtains

$$D_{\text{simple}}^{(1)}(x, y) \simeq -\frac{3}{8\pi^2 r_0^3} \log |x|. \quad (57)$$

However, since $D_{\text{simple}}(x, x)$ grows twice as fast (see Eq. (51)), the combination (54) is bounded in the bulk.

We can consider as well both x and y approaching null infinity in the bulk. In this case, the image y_I approaches the light cone, as illustrated in Fig. 4 (d). Then the logarithm in Eq. (56) is $\sim \log(|x - y_I|/|y_I|^2) \simeq \log(|x||y|)$, hence the combination $\mathcal{G}^{\text{ren}}$ is bounded again.

On the brane, the situation is very different. As mentioned before, $D_{\text{simple}}(x, x)$ is divergent but when properly renormalized, it is simply a constant because of dS symmetry. Then, $\mathcal{G}^{\text{ren}}(x, y)$ behaves like $D_{\text{simple}}(x, y)$ and from Eq. (56) for large separation and $|x| = |y| = r_0$, one obtains

$$\mathcal{G}^{\text{ren}}(x, y) \simeq \frac{3}{4\pi^2 r_0^3} \log |x - y|.$$

Since $|x - y|^2 = 2r_0^2(1 - \cos \zeta)$, this corresponds to the linear growth with the invariant dS time interval $\zeta(x, y)$ between x and y [15].

Finally, we shall mention that the generalization of Eq. (53),

$$\left\langle \left[\frac{\phi(x)}{\mathcal{U}^{\text{bs}}(x)} - \frac{\phi(y)}{\mathcal{U}^{\text{bs}}(y)} \right]^2 \right\rangle^{\text{ren}}$$

⁹ In four dimensions [21], the logarithmic term is $\log |x - y_I|^2/|y_I|^2$. Thus, this string singularity does not appear, despite the same interpretation of Eq. (46) holds.

¹⁰ As an aside, let us comment on the singularities present in the bound state contribution. Specifically, in the term $\partial_p [(p - p_d)G_p^{D-1}]$ present in Eq. (29). We remind that this term is canceled by the KK contribution (28). Hence, it plays no role in the Green’s function. From [14, 15] we know that for x and y timelike related and far apart, then ζ is large and pure imaginary, and it behaves like

$$\partial_p [(p - p_d)G_p^{D-1}] \sim \log |1 - \cos \zeta| = \log \left| \frac{(|x| - |x_I|)^2 - |x - x_I|^2}{2|x||x_I|} \right|.$$

This presents divergences on the light cone and whenever x and y_I (or equivalently, y) are aligned. This condition reduces to the coincidence limit on the brane, but in the bulk one does not expect singularities to appear at all these points. The above mentioned cancelation guarantees that these unphysical singularities are not present in $\mathcal{G}^{\text{ren}}(x, y)$ once we include the KK contribution.

is completely regular in the GK vacuum when $M \neq 0$, and its asymptotic behaviour at infinity parallels that of (53).




FIG. 5: Divergences in $\langle \phi^2 \rangle^{ren} = G(x, x)^{ren}$ and $\langle T_{\mu\nu} \rangle^{ren}$ when the bound state is massless. For the AF vacuum, both the Green's function and the stress tensor diverge at $R = 0$, indicated with a thick dashed line. In addition, if $\xi \neq 0$ the stress tensor diverges on the light cone like $1/U$ or logarithmically in four and six dimensions respectively (thin dashed line). Besides, we also have the ultraviolet ‘Casimir’ divergence on the brane, represented by the plain thick line. In the Garriga Kirsten vacuum (for the massless minimally coupled scalar), $\langle T_{\mu\nu} \rangle^{ren}$ presents only this Casimir-type divergence on the brane.

6. CONCLUSIONS

Our main results can be summarized as follows. In analogy with what happens in de Sitter (dS) space [14, 15], scalar fields with a massless bound state in the spectrum do not have a well defined dS invariant vacuum, except for the massless minimally coupled case. (The case of vanishing bulk mass with non-vanishing curvature coupling has a little subtlety, though.) The Green's function and the v.e.v. of the energy momentum tensor diverges everywhere. The simplest alternative from the analogy to dS case is to take the Allen Follaci vacuum. However, in this vacuum, divergences in the stress tensor are not removed completely within the bulk. Figure 5 illustrates the location of the IR singularities in the Green's function and the stress tensor for the AF vacuum. It remains to clarify whether it is possible or not to avoid these singularities by choosing a vacuum for the KK modes other than the dS invariant vacuum.

When the bound state is very light (but not exactly massless) because M , μ and/or ξ are fine tuned according to Eq. (15), then the stress tensor in dS invariant vacuum takes the form of (24). The stress tensor in this case is smooth, but it becomes very large. Hence, even when the bound state mass is not exactly zero, the dS invariant vacuum looks problematic because of large back reaction. Note that the situation here is different from the usual dS case in two respects. In dS space the large v.e.v. in the stress tensor for the dS invariant vacuum is a constant proportional to the metric. Hence, it might be absorbed by IR renormalization of the cosmological constant. In our case, the stress tensor given by Eq. (24) is a nonlocal expression and cannot be ‘renormalized away’. On the other hand, if one does not want to make any IR renormalization, in the dS case one can take the AF vacuum, and $\langle T_{\mu\nu} \rangle$ stays regular. In the brane world, we do not know the prescription how to remove this large v.e.v. by changing the vacuum state. Choosing non dS invariant vacuum will lead to not only the mentioned divergence in the bulk, but also a new singularity on the light cone, when the bound state mass m_d is not exactly zero. In this case, the radial function for the bound state behaves like $\propto r^{-m_d^2/n}$ near the light cone at $r = 0$. Hence if you single out the contribution to the Green's function from the bound state, it is singular and its derivatives diverges at $r = 0$. Hence, as far as we restrict the change of quantum state to the bound state, we will not be able to remove the large v.e.v. of stress tensor without spoiling its regularity.

A light bound state is compatible with a well behaved and not large stress tensor only in a situation ‘close’ to the massless minimally coupled case. More precisely, here we consider the cases that all of the bulk mass M , the brane mass μ and the nonminimal coupling ξ are small (see Eq. (14)). This corresponds to having a light bound state without accidental cancellations. Namely, the squared bound state mass m_d^2 is of order of the largest among M^2 , $H\mu$

and $H^2\xi$, where H is the Hubble constant on the brane. In this case, the dangerous terms proportional to m_d^{-2} are always associated with some small factor M^2 , μ or ξ , and therefore none of them becomes large.

The case with $M^2 \approx H\mu \ll H^2\xi$ has a little subtlety. In this case, the large v.e.v. in $\langle T_{\mu\nu} \rangle$ appears only in the brane part, and it is constant proportional to the induce metric. Hence, we might be able to consider a model with an appropriate IR renormalization. Only in such a modified model, the stress tensor in the dS invariant vacuum can escape from appearance of a large v.e.v..

Application to the bulk inflaton type models [5, 6] is a part of motivation of the present study. In these models there must be a light bound state of a bulk scalar field. In order to explain the smallness of the bound state mass it will be natural to assume that it is due to smallness of all the bulk and brane parameters without fine tuning. Therefore we will not have to seriously worry about the backreaction of the inflaton in the context of bulk inflaton type models.

We have a few words to add on the massless minimally coupled case. If we consider this case as a limiting situation close to the massless minimally coupled case, the results depend on how we fix the ratio amongst M , $H\mu$ and $H^2\xi$, and hence there remains ambiguity. However, this limiting case has the shift symmetry $\phi \rightarrow \phi + \text{constant}$. If this shift symmetry is one of the symmetries that are to be gauged, there is no ambiguity because the problematic homogeneous mode does not exist in the theory from the beginning. In this setup, undifferentiated ϕ is not an observable. In fact, $\langle T_{\mu\nu} \rangle$ automatically does not contain undifferentiated ϕ . Hence, $\langle T_{\mu\nu} \rangle$ is unambiguously defined although the Green's function is not well defined. We can compute $\langle T_{\mu\nu} \rangle$ in this model by applying the idea of the Garriga Kirsten vacuum (equivalent to a limiting case of AF vacuum), and we confirmed it dS invariant and regular as is expected.

We have also discussed the form of the field perturbations $\langle \phi^2 \rangle$ when the bound state is massless. The main point is that on the brane $\langle \phi^2 \rangle$ grows linearly with dS time χ while in the bulk it is bounded, as expected since the bulk is flat. Aside from this, we derived in closed form the $\langle T_{\mu\nu} \rangle$ for a generic field with zero bulk mass, see Eq. (47).

The same discussion applies in the RSII model with little modifications, which mainly comes from the fact that the Ricci tensor term in the bulk stress tensor $\sim \xi \langle \phi^2 \rangle \mathcal{R}_{\mu\nu}$ is not zero in the RSII model. Thus, the bulk part of $\langle T_{\mu\nu} \rangle$ is finite in the limit of massless bound state only for massless minimal coupling.

Acknowledgments

We are grateful to Jaume Garriga, Misao Sasaki, Wade Naylor, Alan Knapman and Luc Blanchet for useful discussions. T.T. acknowledges support from Monbukagakusho Grant-in-Aid No. 16740141 and Inamori Foundation, and O.P. from JSPS Fellowship number P03193.

APPENDIX A: MASSIVE GREEN'S FUNCTION IN DE SITTER

In this Appendix we obtain the form of the Green's function for a massive field propagating in $D - 1 = n + 1$ dimensional de Sitter (dS) space. The metrics for dS space and its Euclidean version are

$$dS_{(n+1)}^2 = -d\chi^2 + \cosh^2 \chi d\Omega_{(n)}^2 = d\chi_E^2 + \sin^2 \chi_E d\Omega_{(n)}^2, \quad (\text{A1})$$

where $d\Omega_{(n)}^2$ is the metric of a unit n dimensional sphere. The Euclidean time is given by $\chi_E = i\chi + \pi/2$. The Euclidean version of the Hadamard function in dS space can be found from the equation

$$\left[\partial_{\chi_E}^2 + n \cot \chi_E \partial_{\chi_E} - \left(p^2 + \frac{n^2}{4} \right) \right] G_p^{(dS)}(x, x') = -\delta^{(n+1)}(x - x'). \quad (\text{A2})$$

From the symmetry, we can choose x' to be at the pole so that $G_p^{(dS)}$ depends on χ_E only. In terms of $F = (\sin(\chi_E - \pi))^{(n-1)/2} G_p^{(dS)}(x, x')$, this becomes

$$\left[(1 - w^2) \partial_w^2 - 2w \partial_w - \left(p^2 + \frac{1}{4} \right) - \frac{(n-1)^2}{4(1-w^2)} \right] F = 0, \quad (\text{A3})$$

where $w = \cos(\chi_E - \pi)$, and this is solved to give

$$F = N e^{\frac{(n-1)\pi i}{2}} \Gamma\left(\frac{n+1}{2}\right) P_{ip-\frac{1}{2}}^{-\frac{n-1}{2}}(\cos(\chi_E - \pi)). \quad (\text{A4})$$

Hence, the Green's function will be given by the replacement of χ_E by the proper distance between the points x and x' in dS space, which we call $\zeta(x, x')$. $G_p^{(dS)}$ is guaranteed to be regular at $\zeta \rightarrow \pi$. To see the behaviour in the $\zeta \rightarrow 0$ limit, the alternative expression

$$G_p^{(dS)(1)} = \frac{\tilde{N}}{(1 - \cos \zeta)^{\frac{n-1}{2}}} \left\{ F\left(-ip + \frac{1}{2}, ip + \frac{1}{2}, \frac{-n+3}{2}; \frac{1 - \cos \zeta}{2}\right) + \frac{\Gamma(\frac{n}{2} - ip) \Gamma(\frac{n}{2} + ip) \Gamma(-\frac{n-1}{2})}{\Gamma(\frac{1}{2} - ip) \Gamma(\frac{1}{2} + ip) \Gamma(\frac{n-1}{2})} \left(\frac{1 - \cos \zeta}{2}\right)^{\frac{n-1}{2}} F\left(-ip + \frac{n}{2}, ip + \frac{n}{2}, \frac{n+1}{2}; \frac{1 - \cos \zeta}{2}\right) \right\} \quad (A5)$$

is relevant, and here

$$\tilde{N} = \frac{\Gamma(\frac{n+1}{2}) \Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2} - ip) \Gamma(\frac{n}{2} + ip)} N. \quad (A6)$$

In the coincidence limit, the Green's function must behave like $G_{(n+1)}^{(1)} \approx 1/(n-1)S_{(n)}\zeta^{n-1}$, where $S_{(n)}$ is the area of an n -dimensional unit sphere. The limiting behaviour is controlled by the first term. Hence we have $\tilde{N} = 2^{-(n-1)/2}/(n-1)S_{(n)}$.

APPENDIX B: MASSLESS GREEN'S FUNCTION IN DS

Here, we compute the Green's function for the massless scalar in dS. In the massless limit ($p \rightarrow in/2$), the de Sitter (dS) invariant Green's function diverges because of the contribution from the $\ell = 0$ mode, $\mathcal{Y}_{p00}$. The idea is to construct a modified Green's function by substituting this mode by another one that is finite in the limit $p \rightarrow in/2$. In other words,

$$G_{(m=0)}^{(dS)(+)} = G_{(\ell>0)}^{(+)} + \tilde{\mathcal{Y}}\tilde{\mathcal{Y}}'^*,$$

where $\tilde{\mathcal{Y}}$ corresponds to the $\ell = 0$ mode and

$$G_{(\ell>0)}^{(+)} = \sum_{\ell>0} \mathcal{Y}_{p\ell m}(\chi) \mathcal{Y}_{p\ell m}^*(\chi'),$$

where $\mathcal{Y}_{p\ell m}$ are the positive frequency dS invariant vacuum modes. The latter can be obtained as follows. Since $G_p^{(dS)}$ diverges because of the $\ell = 0$ mode, the divergent term in the Laurent expansions of both $G_p^{(dS)}$ and $\mathcal{Y}_{p00}\mathcal{Y}_{p00}'^*$ coincide. We show below that this series contains a simple pole in $p - in/2$. Hence, we can write

$$G_{(\ell>0)}^{(+)} = \lim_{p \rightarrow \frac{in}{2}} \left[G_p^{(dS)(+)} - \mathcal{Y}_{p00}\mathcal{Y}_{p00}'^* \right] = \partial_p \left[\left(p - i\frac{n}{2} \right) G_p^{(dS)(+)} \right]_{p=i\frac{n}{2}} - \partial_p \left[\left(p - i\frac{n}{2} \right) \mathcal{Y}_{p00}\mathcal{Y}_{p00}'^* \right]_{p=i\frac{n}{2}}. \quad (B1)$$

The explicit expression for the first term in the r.h.s. is unnecessary because it is canceled by the KK contribution (28). The dS invariant vacuum mode for $\ell = 0$ with $-ip < n/2$ is

$$\mathcal{Y}_{p00} = C_p \frac{2^{\frac{n-1}{2}} \Gamma(\frac{n+1}{2})}{(\cosh \chi)^{\frac{n-1}{2}}} P_{-ip-\frac{1}{2}}^{-\frac{n-1}{2}}(i \sinh \chi), \quad (B2)$$

and the normalization constant has been chosen so that $\lim_{p \rightarrow in/2} \mathcal{Y}_{p00}/C_p = 1$. The Klein Gordon normalization requires

$$|C_p|^2 = \frac{\Gamma(ip + \frac{n}{2}) \Gamma(-ip + \frac{n}{2})}{2^n \Gamma(\frac{n+1}{2})^2 S_{(n)}}. \quad (B3)$$

Expanding around $p = in/2$ we find

$$|C_p|^2 = \frac{1}{inS_{(n+1)}} \frac{1}{p - in/2} + \mathcal{O}[(p - in/2)^0], \quad (B4)$$

where $S_{(n+1)}$ is the area of an $n+1$ dimensional sphere of unit radius. The behaviour of $\mathcal{Y}_{p00}$ near $p = in/2$, is most easily found using the equation that it solves,

$$\left[\partial_\chi^2 + n \tanh \chi \partial_\chi + \left(p^2 + \frac{n^2}{4} \right) \right] \mathcal{Y}_{p00} = 0. \quad (\text{B5})$$

Bearing in mind that the positive frequency function for the dS invariant vacuum is determined by the regularity when the function is continued to the Euclidean region on the side that contains $\chi = -\pi i/2$, one finds

$$\mathcal{Y}_{p00}(\chi) = C_p \left[1 - \left(p^2 + \frac{n^2}{4} \right) \int^\chi d\chi_1 \frac{1}{\cosh^n \chi_1} \int_{-\pi i/2}^{\chi_1} d\chi_2 \cosh^n \chi_2 + \mathcal{O} \left((p - in/2)^2 \right) \right]. \quad (\text{B6})$$

From this result, the last term in Eq. (B1) can be readily evaluated, and Eq. (32) follows.

APPENDIX C: DIVERGENCE IN THE GREEN'S FUNCTION AND THE $\ell = 0$ MODE

Here, we show that the IR divergence in the Green's function (22) or (44) is due to the homogeneous $\ell = 0$ mode of the bound state. Using Eqs. (11), (21), (B4) and (B6), the contribution from the $\ell = 0$ mode of the bound state for p_d close to $in/2$ is found to be

$$\begin{aligned} G_{\ell=0, p_d}^{(1)} &= \mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r') \mathcal{Y}_{p_d 00}(\chi) \mathcal{Y}_{p_d 00}^*(\chi') + \text{c.c.} \simeq 2 \mathcal{U}^{bs}(r) \mathcal{U}^{bs}(r') |C_{p_d}|^2 [1 + \mathcal{O}(p_d - in/2)] \\ &\simeq \frac{1}{S_{(n+1)}} \frac{1}{p_d - in/2} \frac{I_{n/2}(Mr) I_{n/2}(Mr') / (rr')^{n/2}}{I_{n/2}(Mr_0) \partial_p (\nu_c I_{-ip}(Mr_0) - Mr_0 I'_{-ip}(Mr_0)) \big|_{p=in/2}} + \dots \end{aligned} \quad (\text{C1})$$

where the dots denote higher order in $p_d - in/2$. In the same limit, ν is close to ν_c . The $j = k = 0$ term in (22) is

$$\begin{aligned} G_{j=0, k=0}^{(ren)(1)} &= - \frac{n \Gamma(n)}{2^n \Gamma(\frac{n+1}{2})^2 S_{(n)}} \frac{\nu K_{n/2}(Mr_0) - Mr_0 K'_{n/2}(Mr_0)}{\nu I_{n/2}(Mr_0) - Mr_0 I'_{n/2}(Mr_0)} \frac{I_{n/2}(Mr) I_{n/2}(Mr')}{(rr')^{n/2}} \\ &= - \frac{1}{S_{(n+1)}} \frac{\nu_c K_{n/2}(Mr_0) - Mr_0 K'_{n/2}(Mr_0) + (\nu - \nu_c) K_{n/2}(Mr_0)}{I_{n/2}(Mr_0) (\nu - \nu_c)} \frac{I_{n/2}(Mr) I_{n/2}(Mr')}{(rr')^{n/2}} \\ &\simeq - \frac{1}{S_{(n+1)}} \frac{I_{n/2}(Mr) I_{n/2}(Mr') / (rr')^{n/2}}{I_{n/2}^2(Mr_0) (\nu - \nu_c)} + \mathcal{O}[(\nu - \nu_c)^0] \end{aligned} \quad (\text{C2})$$

where in the second line we used Eq. (15) and the Wronskian relation $K_{n/2}(z) I'_{n/2}(z) - K'_{n/2}(z) I_{n/2}(z) = 1/z$. Using Eq. (12), it is easy to see that

$$\nu - \nu_c = \partial_p \nu \big|_{p_d=in/2} \left(p_d - i \frac{n}{2} \right) + \dots = - \frac{\partial_p [\nu_c I_{-ip} - Mr_0 I'_{-ip}(Mr_0)] \big|_{p=in/2}}{I_{n/2}(Mr_0)} \left(p_d - i \frac{n}{2} \right) + \dots, \quad (\text{C3})$$

so (C1) and (C2) agree in the limit $p_d \rightarrow in/2$. Note that since in this limit this is the dominant contribution and $m_d \simeq in(p_d - in/2)$, the total Green's function (22) can be rewritten in the simple form

$$G_{(ren)}^{(1)}(x, x') \simeq \frac{2}{S_{(n+1)}} \frac{\mathcal{U}_0^{bs}(r) \mathcal{U}_0^{bs}(r')}{r_0^2 m_d^2} + \mathcal{O}(m_d^0), \quad (\text{C4})$$

where $\mathcal{U}_0^{bs}(r) = N_0 I_{n/2}(r)/r^{n/2}$ is the wave function of the bound state for the exactly massless case.

-
- [1] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999) [arXiv:hep-th/9906064].
[2] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221].
[3] W. Naylor and M. Sasaki, Phys. Lett. B **542**, 289 (2002) [arXiv:hep-th/0205277]; E. Elizalde, S. Nojiri, S. D. Odintsov and S. Ogushi, Phys. Rev. D **67**, 063515 (2003) [arXiv:hep-th/0209242]; I. G. Moss, W. Naylor, W. Santiago-German and M. Sasaki, Phys. Rev. D **67**, 125010 (2003) [arXiv:hep-th/0302143]; I. Brevik, K. A. Milton, S. Nojiri and S. D. Odintsov, Nucl. Phys. B **599**, 305 (2001) [arXiv:hep-th/0010205].

- [4] S. Nojiri and S. D. Odintsov, JCAP **0306**, 004 (2003) [arXiv:hep-th/0303011].
- [5] S. Kobayashi, K. Koyama and J. Soda, Phys. Lett. B **501**, 157 (2001) [arXiv:hep-th/0009160].
- [6] Y. Himemoto and M. Sasaki, Phys. Rev. D **63**, 044015 (2001) [arXiv:gr-qc/0010035].
- [7] A. Romeo and A. A. Saharian, J. Phys. A **35**, 1297 (2002) [arXiv:hep-th/0007242].
- [8] A. Knapman and D. J. Toms, Phys. Rev. D **69**, 044023 (2004) [arXiv:hep-th/0309176].
- [9] J. Garriga and M. Sasaki, Phys. Rev. D **62**, 043523 (2000) [arXiv:hep-th/9912118].
- [10] N. Sago, Y. Himemoto and M. Sasaki, Phys. Rev. D **65**, 024014 (2002) [arXiv:gr-qc/0104033].
- [11] Y. Himemoto, T. Tanaka and M. Sasaki, Phys. Rev. D **65**, 104020 (2002) [arXiv:gr-qc/0112027].
- [12] K. Koyama and K. Takahashi, Phys. Rev. D **67**, 103503 (2003) [arXiv:hep-th/0301165];
ibid. D **68**, 103512 (2003) [arXiv:hep-th/0307073].
- [13] N. D. Birrell and P. C. W. Davies, “Quantum Fields In Curved Space,” Cambridge University Press.
- [14] B. Allen and A. Folacci, Phys. Rev. D **35**, 3771 (1987);
A. Folacci, J. Math. Phys. **32**, 2828 (1991) [Erratum-*ibid.* **33**, 1932 (1992)].
- [15] K. Kirsten and J. Garriga, Phys. Rev. D **48**, 567 (1993) [arXiv:gr-qc/9305013].
- [16] A. Vilenkin and L. H. Ford, Phys. Rev. D **26**, 1231 (1982).
- [17] A. D. Linde, Phys. Lett. B **116**, 335 (1982).
- [18] A. A. Starobinsky, Phys. Lett. B **117**, 175 (1982).
- [19] A. Vilenkin, Nucl. Phys. B **226**, 527 (1983).
- [20] S. W. Hawking and I. G. Moss, Nucl. Phys. B **224**, 180 (1983).
- [21] X. Montes, Int. J. Theor. Phys. **38**, 3091 (1999) [arXiv:gr-qc/9904022].
- [22] A. Vilenkin, Phys. Lett. B **133** (1983) 177.
- [23] J. Ipser and P. Sikivie, Phys. Rev. D **30**, 712 (1984).
- [24] A. A. Saharian, Phys. Rev. D **69**, 085005 (2004) [arXiv:hep-th/0308108].
- [25] G. Kennedy, R. Critchley and J. S. Dowker, Annals Phys. **125**, 346 (1980).
- [26] D. Deutsch and P. Candelas, Phys. Rev. D **20**, 3063 (1979).
- [27] S. A. Fulling, J. Phys. A **36**, 6529 (2003) [arXiv:quant-ph/0302117].
- [28] D. Langlois and M. Sasaki, Phys. Rev. D **68**, 064012 (2003) [arXiv:hep-th/0302069].
- [29] J. P. Gazeau, J. Renaud and M. V. Takook, Class. Quant. Grav. **17**, 1415 (2000) [arXiv:gr-qc/9904023].